

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**n artificial intel l igence framewor for compensating transgressions and its appl ication to diet management**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1634422> since 2019-02-06T11:09:) 3Z

*Published version:*

DOI:10.1016/j.jbi.2017.02.015

*Terms of use:*

**Open Access**

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

#### Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.springerlink.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

# Metadata of the article that will be visualized in OnlineFirst

**Please note: Images will appear in color online but will be printed in black and white.**

ArticleTitle	A density management diagram for Norway spruce in the temperate European montane region	
Article Sub-Title		
Article CopyRight	Springer-Verlag Berlin Heidelberg (This will be the copyright line in the final PDF)	
Journal Name	European Journal of Forest Research	
Corresponding Author	Family Name	<b>Vacchiano</b>
	Particle	
	Given Name	<b>Giorgio</b>
	Suffix	
	Division	Department of AGROSELVITER
	Organization	University of Torino
	Address	Via L. da Vinci 44, 10095, Grugliasco, TO, Italy
	Email	gvacchiano@gmail.com
Author	Family Name	<b>Derosé</b>
	Particle	
	Given Name	<b>R. Justin</b>
	Suffix	
	Division	
	Organization	Rocky Mountain Research Station, Forest Inventory and Analysis
	Address	507 25th Street, 84401, Ogden, UT, USA
	Email	
Author	Family Name	<b>Shaw</b>
	Particle	
	Given Name	<b>John D.</b>
	Suffix	
	Division	
	Organization	Rocky Mountain Research Station, Forest Inventory and Analysis
	Address	507 25th Street, 84401, Ogden, UT, USA
	Email	
Author	Family Name	<b>Svoboda</b>
	Particle	
	Given Name	<b>Miroslav</b>
	Suffix	
	Division	Department of Silviculture, Faculty of Forestry and Wood Sciences
	Organization	Czech University of Life Sciences
	Address	Kamycka 129, 16521, Praha 6 Suchdol, Czech Republic
	Email	
Author	Family Name	<b>Motta</b>
	Particle	
	Given Name	<b>Renzo</b>
	Suffix	

Division	Department of AGROSELVITER
Organization	University of Torino
Address	Via L. da Vinci 44, 10095, Grugliasco, TO, Italy
Email	

Schedule	Received	10 August 2012
	Revised	4 February 2013
	Accepted	26 February 2013

Abstract Norway spruce is one of the most important conifer tree species in Europe, paramount for timber provision, habitat, recreation and protection of mountain roads and settlements from natural hazards. Although natural Norway spruce forests can exhibit diverse structures, even-aged stands can arise after disturbance and are the result of common silvicultural practice, including off-site afforestation. Many even-aged Norway spruce forests are actively managed, facing issues such as senescence, insufficient regeneration, mechanical stability (stem form), sensitivity to biotic disturbances, and restoration. We propose the use of Density Management Diagrams (DMD), stand-scale graphical models originally designed to project growth and yield of even-aged forests, as a heuristic tool for assessing the structure and development of even-aged Norway spruce stands. DMDs are predicated on basic tree allometry and the assumption that self-thinning occurs predictably in forest stands. We designed a DMD for Norway spruce in temperate Europe based on wide-ranging forest inventory data. Quantitative relationships between tree- and stand-level variables that describe resistance to selected natural disturbances were superimposed on the DMD. These susceptibility zones were used to demonstrate assessment and possible management actions related to, for example, windfirmness and effectiveness of the protective function against rockfall or avalanches. The Norway spruce DMD provides forest managers and silviculturists a simple, easy-to-use, tool for evaluating stand dynamics and scheduling needed density management actions.

Keywords (separated by '-')	Decision support systems - Natural hazards - <i>Picea abies</i> (L.) Karst. - Protective function - Self-thinning - Silviculture
-----------------------------	----------------------------------------------------------------------------------------------------------------------------------

Footnote Information	Communicated by A. Weiskittel.
----------------------	--------------------------------

Journal: 10342  
Article: 694



## Author Query Form

**Please ensure you fill out your response to the queries raised below  
and return this form along with your corrections**

Dear Author

During the process of typesetting your article, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the 'Author's response' area provided below

Query	Details required	Author's response
1.	Please check whether the mail ID "giorgio.vacchiano@unito.it" should appear in the publication.	
2.	Please check and confirm the city name is correctly identified in the affiliation part.	
3.	References Hynynen (1993), R Development Core Team (2011) are given in list but not cited in text. Please cite in text or delete from list.	
4.	Please check and confirm the journal title of the reference Baier (1996).	
5.	Reference Hynynen and Ojansuu (2003) is given in list but not cited in text. Please cite in text or delete from list.	

# A density management diagram for Norway spruce in the temperate European montane region

Giorgio Vacchiano · R. Justin Derose ·  
John D. Shaw · Miroslav Svoboda ·  
Renzo Motta

Received: 10 August 2012 / Revised: 4 February 2013 / Accepted: 26 February 2013  
© Springer-Verlag Berlin Heidelberg 2013

**Abstract** Norway spruce is one of the most important conifer tree species in Europe, paramount for timber provision, habitat, recreation and protection of mountain roads and settlements from natural hazards. Although natural Norway spruce forests can exhibit diverse structures, even-aged stands can arise after disturbance and are the result of common silvicultural practice, including off-site afforestation. Many even-aged Norway spruce forests are actively managed, facing issues such as senescence, insufficient regeneration, mechanical stability (stem form), sensitivity to biotic disturbances, and restoration. We propose the use of Density Management Diagrams (DMD), stand-scale graphical models originally designed to project growth and yield of even-aged forests, as a heuristic tool for assessing the structure and development of even-aged Norway spruce stands. DMDs are predicated on basic tree allometry and the assumption that self-thinning occurs predictably in forest stands. We designed a DMD for Norway spruce in temperate Europe based on wide-ranging forest inventory data. Quantitative relationships between tree- and stand-level variables that describe resistance to selected natural

disturbances were superimposed on the DMD. These susceptibility zones were used to demonstrate assessment and possible management actions related to, for example, windfirmness and effectiveness of the protective function against rockfall or avalanches. The Norway spruce DMD provides forest managers and silviculturists a simple, easy-to-use, tool for evaluating stand dynamics and scheduling needed density management actions.

**Keywords** Decision support systems · Natural hazards · *Picea abies* (L.) Karst. · Protective function · Self-thinning · Silviculture

## Introduction

Norway spruce (*Picea abies* (L.) Karst.) is one of the most important tree species in the mountain ranges of central and southern Europe. Norway spruce stands are important for timber production and provide important ecosystem services (Pretzsch et al. 2008). In mountain regions, these forests can provide protection from natural hazards such as avalanches, rockfall, or landslides (Bebi et al. 2001; Mayer and Ott 1991). Norway spruce forests also provide habitat for game, and may harbor endangered fauna or flora (e.g., Nascimbene et al. 2009).

Vast areas of pure, monolayered Norway spruce plantations are common in many European montane and lowland landscapes, oftentimes usurping the space of natural forests (Hansen and Spiecker 2004). The species has been introduced far outside its natural range, both in countries where it occurs naturally, for example, in Germany and Norway, and in novel areas such as Denmark, Belgium, and Ireland (Skroppa 2003). Natural and semi-natural Norway spruce forests, on the other hand, are relatively

Communicated by A. Weiskittel.

G. Vacchiano (✉) · R. Motta  
Department of AGROSELVITER, University of Torino,  
Via L. da Vinci 44, 10095 Grugliasco, TO, Italy  
e-mail: gvacchiano@gmail.com

R. J. Derose · J. D. Shaw  
Rocky Mountain Research Station, Forest Inventory  
and Analysis, 507 25th Street, Ogden, UT 84401, USA

M. Svoboda  
Department of Silviculture, Faculty of Forestry and Wood  
Sciences, Czech University of Life Sciences, Kamycka 129,  
Praha 6 Suchbát 16521, Czech Republic

rare (Parviainen et al. 2000; Motta 2002) and often exhibit multiple structural and compositional attributes depending in part on the disturbance regime (Shohorova et al. 2009). These structures range from sparse, multilayered subalpine stands (Kulakowski et al. 2004; Krumm et al. 2011) to monolayered forests resulting from severe disturbances (Fisher et al. 2002; Angelstam and Kuuluvainen 2004), to uneven-aged mixtures (Svoboda et al. 2010, 2012).

Windstorms, snow loading, and insects are among the most damaging disturbance agents in Norway spruce stands (Klopčič et al. 2009; Svoboda et al. 2012). Increasing susceptibility to natural disturbances (Schlyter et al. 2006; Seidl et al. 2011), in combination with aging stands and increasing demand for enhanced structural complexity and close-to-nature forest structures (Gamborg and Larsen 2003), results in a silvicultural conundrum that cannot be adequately addressed using simple management tools (e.g., yield tables). Given the importance of Norway spruce in managed montane forests of central-southern Europe, it is important to develop ecologically based decision support systems that allow for the development of realistic management scenarios and enable the comparison of alternative schedules with respect to the evaluation criteria of interest (e.g., volume production, carbon storage, stand stability, structural diversity, nature conservation, and biodiversity).

Density management diagrams (DMD) are empirical models of even-aged stand dynamics (Jack and Long 1996). They reflect fundamental relationships involving tree size, stand density, site occupancy, and self-thinning. Allometric relationships between mean tree size, age, height, and yield are portrayed allowing users to design treatments by plotting both current and desired future stand structure on the DMD. Alternative management strategies that accomplish diverse objectives can be simultaneously compared and their efficacy evaluated at a glance. In this paper, we analyzed data from Norway spruce stands to construct a DMD with wide applicability across montane regions of central-southern Europe. Using specific examples of (1) maximizing volume production, (2) mechanical stability against wind damage, (3) avalanche protective function, and (4) potential resistance to spruce bark beetle (*Ips typographus* L.), we demonstrate the usefulness of the Norway spruce DMD.

## Methods

### Data sources

The data used to develop the Norway spruce DMD came from multiple sources (Table 1) that covered many regions of central-southern Europe (Fig. 1) and included 5,656


plots. Most areas occupied by temperate European montane forest were represented in the data set. We excluded areas with few pure Norway spruce forests (e.g., Balkans) or countries where forest inventory data were not readily accessible.

1. Data from France were obtained from the French National Forest Inventory (<http://www.ifn.fr/spip/>) for the inventory period 2005–2009. The French inventory design implemented three nested fixed-area plots [6, 9, and 15 m radius for trees ~7–22.5 cm, 22.6–37.5 cm, and 37.5+ cm in diameter at breast height (DBH), respectively] from which trees per hectare (N) expansion factors were calculated. The French Inventory also included tree height (H) and estimated tree volume (Vidal et al. 2007).
2. Data from the Czech Republic came from two regions, Sumava and Tajga. In the Sumava region, the inventory design was three nested fixed-area plots (3.5, 7, and 12.6 m radius for trees 7–14.9 cm, 15–29.9, and 30+ cm DBH, respectively) and did not include estimates of tree volumes (Čížková et al. 2011). In the Tajga region, the inventory consisted of one 12.5-m-radius fixed-area plot where DBH and H were measured and estimates of volume included for all trees >10 cm.
3. Data from Romania came from the mountain regions of Călimani and Giumalău (Cenușă 1992). The inventory in these regions used either a 500- or 1,000-m<sup>2</sup> fixed-area plot with a lower DBH cutoff of 10 cm. Individual-tree heights for all trees were estimated using locally calibrated models and there were no estimates of volume (M. Svoboda—unpublished data).
4. Italian data came from multiple regions and inventory designs. At Aosta and Piemonte (IPLA 2003) fixed-area plots ranged from 8 to 15 m radius depending on overstory density and the lower DBH cutoff was ~7 cm; species- and site-specific volume equations were provided. At Paneveggio and San Martino (Berretti and Motta 2005) fixed-area plots of 12 m radius with a lower DBH cutoff of 17 cm were used and no estimates of volume were made. At Val Pontebbana (Castagneri et al. 2010) 12-m-radius fixed-area plots were sampled with a lower DBH cutoff of ~7 cm. In Valbona, 400-m<sup>2</sup> fixed-area plots were used with a lower DBH cutoff of ~7 cm (Motta et al. 2006). At Burgusio, Lasa, Latemar, Luttago, Meltina, Naturno, Valle Aurina, and for plots of the National Forest Inventory (INFC 2006), variable radius plots (basal area factor = 4 m<sup>2</sup> ha<sup>-1</sup>) were employed with a lower DBH cutoff of ~4 cm and volume was not estimated.

**Table 1** Source of data for the Norway spruce DMD and estimates of  $SDI_{max}$  by location ( $SDI_p$  for pure, even-aged Norway spruce stands)

ID	Dataset name (region)	Country	No. of plots	DBH cutoff (cm)	Plot size (m <sup>2</sup> )	98th percentile $SDI_{max}$
1	Aosta	Italy	156	7	201–707	1,209
2	Piemonte	Italy	65	7	201–707	1,701
3	National forest inventory	Italy	401	4	Relascope	1,571
4	Burgusio	Italy	91	4	Relascope	1,080
5	Lasa	Italy	251	4	Relascope	1,473
6	Latemar	Italy	322	4	Relascope	1,745
7	Luttago	Italy	72	4	Relascope	1,007
8	Meltina	Italy	256	4	Relascope	1,383
9	Naturno	Italy	304	4	Relascope	1,220
10	Valle Aurina	Italy	155	4	Relascope	1,493
11	Paneveggio	Italy	91	17	452	1,321
12	San Martino	Italy	91	17	452	1,278
13	Valbona	Italy	66	7	400	1,592
14	Val Pontebbana	Italy	33	7	452	1,162
15	Tajga	Czech Republic	78	7	491	755
16	Sumava Certovo	Czech Republic	66	7	38–499	1,278
17	Sumava NP	Czech Republic	38	7	38–499	1,221
18	Sumava large plots	Czech Republic	15	7	1,000–2,500	1,121
19	Sumava Trojmezna	Czech Republic	18	7	38–499	826
20	Călimani	Romania	40	10	500–1,000	1,425
21	Giulmalau	Romania	41	10	500–1,000	1,270
22	Baden-Württemberg	Germany	399	7	Relascope	1,464
23	France 2005	France	522	7	113–707	1,206
24	France 2006	France	526	7	113–707	1,277
25	France 2007	France	558	7	113–707	1,305
26	France 2008	France	489	7	113–707	1,086
27	France 2009	France	471	7	113–707	1,238
28	Parangalitsa	Bulgaria	227	4	100	2,653

**Fig. 1** Distribution of Norway spruce in central-southern Europe (after Schmidt-Vogt 1977) and location code for data used for DMD construction. Refer to Table 1 for location names





- 162 5. Bulgarian data referred to remote-sensed, internally  
163 homogenous forest patches in the Parangalitsa Reserve,  
164 including a number of post-disturbance stands (Panayotov  
165 et al. 2011). A total of 227 100-m<sup>2</sup> plots were sampled with  
166 a lower DBH cutoff of 4 cm and no information on  $H$  and  
167 volume.  
168 6. German data came from the Second National Forest  
169 Inventory of Germany (Schmidt and Kandler 2009).  
170 Trees with a minimum DBH of 7 cm were selected  
171 using the angle-count method (horizontal point sam-  
172 pling) with a basal area factor of 4 m<sup>2</sup> ha<sup>-1</sup>. The  
173 attributes recorded included species, DBH, tree age,  
174 and  $H$ .

## 175 Size–density relationships

176 Using the tree-level data, we calculated  $N$ , quadratic mean  
177 diameter (QMD), basal area, percent basal area of Norway  
178 spruce, stand density index (SDI), and stand top height  
179 (HT<sub>100</sub>), defined as the average height of the 100 largest  
180 (DBH) trees per hectare. SDI was calculated two ways: (1)  
181 Reineke (SDI<sub>p</sub>: Reineke 1933, modified by Long and  
182 Daniel 1990),

$$SDI_p = N(QMD/25.4)^{1.605}, \quad (1)$$

184 and (2) summing the SDI of each  $i$  th tree in a stand  
185 (SDI<sub>sum</sub>: Shaw 2000),

$$SDI_{sum} = \sum N_i (DBH_i/25.4)^{1.605} \quad (2)$$

187 so that stands with simple structure could be filtered from  
188 the data using the SDI<sub>sum</sub>:SDI<sub>p</sub> ratio (SDI<sub>ratio</sub>). SDI<sub>ratio</sub> has  
189 been shown to theoretically differentiate even-aged stands,  
190 which have strong unimodal diameter distributions (SDI<sub>r-  
191 ratio</sub> ≥ 0.9), from uneven-or multi-aged stands, which show  
192 increasing skewness in their diameter distribution (SDI<sub>r-  
193 ratio</sub> < 0.9) (Ducey 2009). SDI<sub>ratio</sub> has been used to indicate  
194 relatively even-aged stands for building DMDs (Long and  
195 Shaw 2005, Shaw and Long 2007). Before estimating the  
196 self-thinning boundary, the plot-level data were filtered for  
197 Norway spruce composition ≥ 80 % (determined by per-  
198 cent basal area) and for even-aged stands (SDI<sub>ratio</sub> ≥ 0.9),  
199 which resulted in 1,609 plots.

200 We paid particular attention to determining the maxi-  
201 mum size–density line. In order to filter for fully stocked  
202 stands, we used a binning method (Bi and Turvey 1997)  
203 (200 N bins) from which maximum observations of SDI<sub>sum</sub>  
204 were extracted before the maximum self-thinning line was  
205 fit by ordinary least-squares (OLS) regression. We assessed  
206 whether a lower DBH cutoff of 4, 7, 10, or 17 cm had any  
207 effect on SDI<sub>max</sub> (Curtis 2010) and/or the slope determined  
208 during the binning method by refitting the OLS for each

DBH cutoff group. Moreover, since differing self-thinning  
slopes are reported in the literature, both between- and  
within-tree species (including Norway spruce: Sterba 1987;  
Hynynen 1993; Monserud et al. 2005; Pretzsch and Biber  
2005; Pretzsch 2006; Schütz and Zingg 2010; Charru et al.  
2011), we tested whether Reineke's (1933) suggested slope  
of −1.605 was statistically different from that of our linear  
fit. Subsequently, we shifted the OLS line to cross the point  
of maximum stocking. SDI<sub>max</sub> indicates maximum growing  
space occupancy (Yoda et al. 1963), so that plots falling  
above the line should be exceedingly rare. Therefore, we  
assumed the 98th percentile of the SDI<sub>sum</sub> frequency dis-  
tribution appropriately characterized the maximum attain-  
able SDI. Finally, we juxtaposed lines on the DMD to  
describe relative stand density (percent of SDI<sub>max</sub>) fol-  
lowing the recommendations of Long (1985). That is, 25 %  
of SDI<sub>max</sub> represents crown closure, 35 % of SDI<sub>max</sub>  
indicates the beginning of individual-tree growth reduction  
due to inter-tree competition, and at 60 % of SDI<sub>max</sub>, the  
onset of severe competition.

We tested for the existence of a Mature Stand Boundary  
(MSB) in the maximum self-thinning limit (Shaw and  
Long 2007) by fitting the following three-parameter  
function:

$$QMD = a(N_{max} + b)^c, \quad (3)$$

where  $N_{max}$  are observations of maximum  $N$  for each 0.01  
class of Log<sub>10</sub> QMD. Only plots where QMD ≥ 15 cm  
were used, because stands in the smaller size classes are  
not needed to establish the MSB. Subsequently, we shifted  
the curve developed in Eq. 3 such that the maximum SDI  
value on the curve was asymptotic to the SDI<sub>max</sub> on the  
DMD.

## Top height and volume

When included on a DMD, HT<sub>100</sub> can be used with local  
site index curves to assess and quantify the temporal  
development of a particular stand (Jack and Long 1996).  
Using plot data that included observations of HT<sub>100</sub>, we  
modeled QMD as a function of HT<sub>100</sub>, attenuated by an  
inverse logarithmic function of tree density:

$$QMD = HT_{100}(b_1 - b_2 \ln N) \quad (4)$$

To generate stand-level volume (VOL) isolines on the  
DMD, we modeled VOL as a power function of QMD and  
 $N$  (Eq. 5a), then rewrote the equation as QMD =  $f(VOL)$ ,  
where VOL is total standing volume (m<sup>3</sup> ha<sup>-1</sup>) for plot  
data with volume observations:

$$VOL = c_1 + c_2 N QMD^{c_3} \quad (5a)$$

$$QMD = [(c_1 + c_2 N)^{-1} VOL]^{(1/c_3)} \quad (5b)$$

We plotted  $HT_{100}$  and VOL isolines on the DMD for ranges of 20–50 m, and 200–1,200  $m^3 ha^{-1}$ , respectively. Different inventories may have used different equations for tree or stand volume, generating idiosyncrasies when pooling all volume data in one model. However, because we were missing inventory-specific volume equations, we used original data as much as possible, acknowledging that DMD isolines merely represent average conditions across the entire dataset.

All models were assessed for parameter significance and goodness-of-fit by computing adjusted  $R^2$  and root mean square error (RMSE). We determined that both models had little or no bias by inspecting residual plots over the predictor variables, elevation when available, SDI, basal area, region, and whether the plot had a lower DBH cutoff of 4, 7, 10, or 17 cm.

### Disturbances and site index

To illustrate the advantages of the DMD in designing silvicultural strategies to maximize resistance to disturbances and protection from natural hazards, we superimposed “susceptibility zones” on the diagram, which encapsulate combinations of size and density that (a) fulfill an effective protection against avalanche release; and (b) result in a low risk of wind damage. Thresholds for (a) were summarized as follows (after Berretti et al. 2006; Gauquelin and Courbaud 2006):

- Basal area  $\geq 25 m^2 ha^{-1}$  when QMD = 25 cm, and  $\geq 7.5 m^2 ha^{-1}$  when QMD = 10 cm for effective snowpack stabilization if slope is steeper than 35°;
- Live crown ratio  $\geq 60\%$  in trees or cluster of trees supporting the stability of the stand. We relaxed this requirement to  $\geq 33\%$ , representing a minimal acceptable level of individual-tree vigor that should be ensured with a relative SDI  $< 0.60$  (Long 1985);
- H/DBH ratio  $< 80$  in dominant trees. H/DBH (mean or dominant) ratio cannot be read directly off the DMD. However, assuming that DBH is normally distributed in a stand and that dominant diameter (DD) is equivalent to the 90th percentile of such distribution (Z value = +1.64), DD can be computed by

$$DD = 1.64\sigma_{DBH} + QMD, \quad (6a)$$

where  $\sigma_{DBH}$  is the standard deviation of the DBH distribution in the stand. In order to represent risk zones on the DMD, we assumed that  $\sigma_{DBH} = 0.3$  QMD and solved Eq. 6a for QMD:

$$QMD = 0.67 DD, \quad (6b)$$

to be substituted in  $HT_{100}/QMD$  ratio from Eq. 4 and constrained to  $\leq 0.8$ . This allowed the influence of smaller, suppressed trees to be removed so that only the slenderness of dominant trees was considered (Castedo-Dorado et al. 2009);

- Gap size  $\leq 1.5$  times tree height (in order to avoid tree-free patches prone to dangerous snow gliding). If square spacing is assumed, a Mean nearest neighbor distance ( $m$ ) (MNND) can be computed as the square root of the reciprocal of  $N$ . We introduced a multiplier to account for clumped patterns, that is, the ratio between maximum and observed nearest neighbor index (NNI). NNI ranges from 0, when trees are highly clumped, to 2.1491, when trees are arranged along a hexagonal grid (Clark and Evans 1954):


$$MNND = (2.1491 NNI^{-1}) \times 100 N^{-0.5}, \quad (7)$$

subsequently constrained to  $\leq 1.5HT_{100}$  and used to back calculate critical  $N-HT_{100}$  combinations.

While the DMD can be used to assess avalanche hazard related to stand structure, other predisposing conditions (e.g., weather, topography, characteristics of snowpack, and terrain ruggedness) must be evaluated independently.

Thresholds for live crown ratio followed those by Riou-Nivert (2001), who established low, medium, and high wind risk zones for conifer species, based on the relationship between QMD and  $HT_{100}$  (Fig. 2). Mitchell (2000) suggested that such general zones of stability exist for uniform stands of temperate zone conifers.

An appropriate site index (SI) curve allows the estimates of  $HT_{100}$  on the DMD to be a surrogate for time (Drew and Flewelling 1979). SI estimates were not included in the raw data. In order to provide SI curves applicable to even-aged, pure Norway spruce stands across temperate Europe, we fitted a modified Richards' model of height growth (Sterba 1976) to yield tables from Eisacktal, South Tyrol (Moser 1991), which exhibited a wide range of fertility classes (i.e.,  $HT_{100}$ : 7.9–45.8 m at age 100). All statistics were performed in the R environment version 2.14.1 (R Development Core Team 2011).



**Fig. 2** Wind stability zones for even-aged coniferous stands based upon  $HT_{100}$  and QMD (after Riou-Nivert Riou-Nivert 2001)



## Results

Twenty-nine percent of the original Norway spruce data set, that is, 1,609 of 5,656 inventory plots (Table 2) were used to fit a maximum size–density relationship characterizing montane Norway spruce in central-southern Europe. Slope of the self-thinning line was  $-1.497$  (adjusted  $R^2 = 0.94$ ); the 95 % confidence interval of the slope coefficient from OLS regression ( $-1.671$  to  $-1.324$ ) included Reineke's value of  $-1.605$ .  $SDI_{max}$  was 1,461 (Fig. 3); coefficient of variation between the 28 regions was 26 %, mean = 1334.28, and  $SD = 345.39$  (Table 1). Binning by different DBH cutoff values did not change our results with respect to the significance of  $-1.605$ , except for the 17 cm cutoff that produced a non-significant regression slope likely due to limited sample size (Table 3). However, the lowest DBH cutoff (4 cm) produced the highest  $SDI_{max}$ . Parameters of the MSB (Eq. 3) were  $a = 3330.105$ ,  $b = 185.158$ , and  $c = -0.0656$  (adjusted  $R^2 = 0.96$ ).

Top height and volume equations were statistically significant (Table 4). Some bias was revealed in residual plots over observed volume (Fig. 4); however, these occurred in poorly stocked stands (i.e.,  $<50 \text{ m}^3 \text{ ha}^{-1}$ ) and do not constitute a concern for using the DMD in practice. The QMD- $HT_{100}$  model exhibited some high regional bias (Table 5); a 95 % confidence envelope about the mean of QMD residuals included zero in 7 out of 14 sites for the  $HT_{100}$  model (Eq. 4), and 8 out of 10 sites for the VOL model (Eq. 5b).

## Discussion

### DMD characteristics

DMDs that cover widely distributed species (e.g., Long and Shaw 2005) are indicative of average growth patterns

and allometric relationships of monospecific stands. We assumed that allometric equations, when portrayed on the DMD, were invariant across all sites (Weiner 2004). Conditions under which the self-thinning boundary may shift include, at the local scale, genetic differences (Buford and Burkhardt 1987), and severe resource deficiencies, e.g., in tree line environments (Körner 2003). However, despite deviations at certain localities (Table 5), our allometric models should be robust, in that the high number of plots used for calibration should average out local peculiarities.


Previous research has observed disparities in mortality rates of Norway spruce stands located on different elevations and aspects (Krumm et al. 2012). However, we consider these to be an effect of the different rates at which stands may progress along their trajectories of development in size–density space, while following the same overarching, species-specific, self-thinning boundary. Differences in topography, temperature, light, and soil fertility affect growth rates and, in turn, the rate of mortality during the stem-exclusion phase (Aulitzky 1984; Schönenberger 2001). In other words, a Norway spruce stand on a high-quality site will reach the boundary more quickly than the same density of trees on a lower-quality site, even though both eventually achieve the same boundary (Jack and Long 1996). This constancy is fundamental to the general utility of DMD and allows the use of site index curves to determine the time required to attain particular stand structural characteristics. Our aim was to characterize Norway spruce stands across the montane forest region in central-southern Europe using a single tool. Therefore, when using the DMD to portray stands at a specific location, managers should choose the appropriate dominant height curve, in order to account for differences in local productivity.

Maximum  $SDI$  for Norway spruce in montane forests of central-southern Europe was 1,461, which was intermediate in the range of previous regional estimates (Pretzsch 2005—Germany:  $SDI_{max} = 1,609$ ; Monserud et al. 2005—Austria:  $SDI_{max} = 1,571$ ; Sterba 1981—Austria:  $SDI_{max} = 1,547$ ; Castagneri et al. 2008—NE Italy:  $SDI_{max} = 1,380$ ), independent of the DBH measurement cutoff. Consistent with previous studies (Shaw and Long 2007), we detected a convex pattern to the self-thinning limit at high tree size–low density combinations, that is, a mature stand boundary (MSB). The most commonly suggested explanation for this process is so-called “self-tolerance” (Zeide 1985), by which growing space resulting from the death of very large trees can not be promptly reclaimed by con-specific neighboring trees, lowering the limit of possible size–density combinations. Maintaining stand size–density below the MSB is crucial for management as combinations above the line are ecologically improbable (DeRose et al. 2008).

**Table 2** Summary statistics for pure, even-aged Norway spruce stands ( $SDI_{ratio} \geq 0.9$ , percent Norway spruce on total basal area  $\geq 0.8$ )

Variable	Unit	<i>n</i>	Min	Max	Mean	SE
N	trees $\text{ha}^{-1}$	1,609	14	5,058	564.1	13.03
QMD	cm	1,609	7.8	115.0	34.8	0.31
$HT_{100}$	m	876	4.2	46.0	24.1	0.23
VOL	$\text{m}^3 \text{ ha}^{-1}$	505	0.8	1163.6	316.4	9.69
BA	$\text{m}^2 \text{ ha}^{-1}$	1,609	0.4	130.0	40.3	0.50
PRCPA	%	1,609	0.8	1.0	1.0	0.002
$SDI_{sum}$	—	1,609	14	2,057	705.0	8.45
$SDI_{ratio}$	—	1,609	0.9	1.0	1.0	0.001
Age	Years	669	8.0	338.0	108.5	2.40
Elevation	m. a.s.l.	748	82	2,230	1240.6	16.26

**Fig. 3** Selected Norway spruce stands in size–density space, SDI lines, and mature stand boundary



**Table 3** Fit statistics of the self-thinning line computed using different DBH cutoff values

DBH cutoff (in.)	SDI <sub>max</sub>	Slope	95 % min	95 % max	<i>p</i>	Adjusted <i>R</i> <sup>2</sup>	No. of plots
0	1,461	−1.50	−1.67	−1.32	0.00	0.94	1,609
4	1,587	−1.61	−1.85	−1.36	0.00	0.90	633
7.5	1,287	−1.53	−1.95	−1.10	0.00	0.82	635
10	1,447	−1.52	−1.83	−1.20	0.00	0.91	250
17	1,355	−1.87	−3.77	0.04	0.053	0.56	91

Application of the DMD


The DMD is depicted in log(QMD)-log(Density) space with a superimposed self-thinning line and HT<sub>100</sub> and VOL isolines (Fig. 5). Application of the DMD proceeds as follows: (1) identify starting conditions on the DMD (i.e., current stand structure); (2) identify target stand structure at

end of rotation (EOR) and track the likely trajectory of unmanaged stand development (i.e., asymptotic to the self-thinning boundary); (3) ascertain the need for stand density regulation, e.g., to prevent the onset of competition-related mortality (~ 60 % SDI) and represent the planned thinning entries on the DMD; (4) assess time to reach EOR by tracking the starting and ending HT<sub>100</sub> on SI curves (Fig. 6).



**Table 4** Model fit and parameters for Eqs. 4 and 5b (HT<sub>100</sub> in m, QMD in cm, VOL in m<sup>3</sup> ha<sup>-1</sup>)

Parameter	Estimate	S.E.	95 % min	95 % max	Adjusted $R^2$	$n$
QMD = HT <sub>100</sub> ( $b_1 - b_2 \ln N$ )						
$b_1$	3.148	0.056	3.038	3.259	0.663	1,491
$b_2$	0.297	0.009	0.278	0.315		
VOL = $c_1 + c_2 N$ QMD <sup><math>c_3</math></sup>						
$c_1$	-25.795	5.238	-36.087	-15.503	0.937	505
$c_2$	$1.79 \times 10^{-4}$	$1.6 \times 10^{-5}$	$1.46 \times 10^{-4}$	$2.11 \times 10^{-4}$		
$c_3$	2.432	0.025	2.383	2.480		



**Fig. 4** Residual plots from HT<sub>100</sub> (a) and VOL (b) models (Eqs. 4 and 5b). Black lines represent loess fit

*Maximize volume production*

When the goal is timber production, one can use the DMD for minimizing the time required to reach EOR at a desired mean stem diameter. In addition, by using the HT isolines in combination with site-specific potential productivity, one can incorporate future revenue and future costs into the density management regime. For example, if the desired EOR QMD was 40 cm, and the current stand has ~2,600 N (see Fig. 5), a thinning would be necessary to forestall density-dependent mortality when relative SDI approaches 60 %. This could be achieved by pre-commercially thinning the stand to ~400 N. This would drive stand development on a trajectory to meet the desired EOR of 40 cm at approximately the same time maximum stand growth is achieved (relative SDI = 60 %). Both the timing

and volume of the pre-commercial thinning, or any subsequent commercial thinnings could be estimated using the HT and VOL isolines, respectively, and the return or cost associated with that treatment discounted to today's values to compare management alternatives. Similar to a volume-based regime, by using appropriate biomass conversion factors, and assuming a carbon conversion factor of 0.5, one could plan a density management regime to maximize aboveground carbon sequestration for a particular stand.


*Mechanical stability against wind damage*


Windstorms are the most destructive disturbance agent in temperate European forests (judged by the volume of timber damaged: Schelhaas et al. 2003), often causing extensive damage in Norway spruce, and in particular in

**Table 5** QMD mean bias (predicted–observed 95 % confidence interval) for HT<sub>100</sub> and VOL models (Eqs. 4 and 5b), by location

Location	Mean bias QMD–HT <sub>100</sub> (cm)		Mean bias QMD–VOL (cm)	
	Lower	Upper	Lower	Upper
Aosta	1.31	3.17	–0.14	0.37
Piemonte	–5.49	1.88	–1.82	–0.37
Italy	–0.30	1.33	–	–
Valbona	–3.67	–1.23	–0.39	1.00
Val Pontebbana	–3.52	0.68	–0.62	1.05
Tajga	2.34	3.57	0.74	2.82
Sumava NP	–2.99	–1.10	–	–
Călimani	3.29	5.22	–	–
Giumalau	5.41	8.14	–	–
France 2005	–3.57	–0.13	–0.43	1.78
France 2006	–3.05	–0.47	–0.52	1.06
France 2007	–3.46	0.77	–0.48	2.24
France 2008	–1.22	2.80	–0.04	2.67
France 2009	–2.51	0.50	–0.76	0.89

**Fig. 5** DMD for Norway spruce in the central-southern European montane ecoregion, and working example of stand trajectories for unmanaged and a pre-commercial thinning alternative (starting stand conditions:  $N = 2,500$ , QMD = 10 cm; end of rotation: QMD = 40 cm). Competition-related mortality onsets at 60 % SDI and higher. Target QMD is reached in 70 years in the working example, as opposed to 90 years in the unmanaged alternative, on a medium fertility site ( $SI = 23.6$  m, see Fig. 6)





**Fig. 6** Site index curves from Eisacktal (South Tyrol) yield tables

structurally homogeneous stands (Schmidt-Vogt et al. 1987). Tree damage begins at wind speeds of  $15 \text{ m s}^{-1}$  and can be catastrophic at  $25 \text{ m s}^{-1}$  (Zajackowski 1991). Susceptibility is higher for slender trees (e.g., Rottmann 1986; Thomasius 1988; Riou-Nivert 2001; Dobbertin 2002) and short, broad crowns (Schütz et al. 2006), a condition created through stand dynamics characterized by intense inter-tree competition. When risk zones for wind damage are superimposed on the DMD (Fig. 7), two types of management action are supported: (1) the ability to assess current conditions relative to risk, and (2) the possibility of projecting the effect of interventions which aim to maintain or drive stand structures into low-risk areas as long or quickly as possible. For example, the second management approach is depicted in the example of an unmanaged stand trajectory portrayed in Fig. 7. Among structural attributes, a threshold of  $\sim 1,800 \text{ trees ha}^{-1}$  strongly differentiates high and medium susceptibility to wind damage. By contrast, the threshold-to-low susceptibility is mainly determined by tree slenderness, where “safe” values are typically encountered in low-density stands. From such results, we conclude that the typical even-aged Norway spruce stand (either natural or planted) is characterized by a medium risk of wind damage.

First glance at our Norway spruce stand plotted on the DMD might indicate that a heavy thinning may effectively lower stand susceptibility to wind damage, but in dense stands, it may result in sudden isolation of trees with high


height-to-diameter ratio, and hence, increase the probability of damage by breakage or uprooting (Thomasius 1980).

While uneven-aged stands are acknowledged to have higher resistance to wind (e.g., Shorohova et al. 2008), they cannot be accurately represented on the diagram. Additional limitations of DMD are (a) they cannot track risk factors unrelated to stand structure, for example, soil (trees are much more vulnerable to wind damage on shallow or wet soils), weather, and building beetle populations; and (b) they cannot track the long-term influence of climate change on either autogenic, or allogenic growth factors.

#### *Avalanche and rockfall protective function*

Because Norway spruce predominates in the upper montane and subalpine belt, it can be quite effective against the release of avalanches (although not on their transit), provided that stands meet given structure and density standards (Motta and Haudemand 2000). Like wind damage, required stand structures can be represented as risk zones on the DMD (Fig. 8). Although individual-tree resistance parameters are similar to those required for windfirmness, effective stand structures differ because open stands with thicker trees are more prone to avalanche release due to the presence of tree-free gaps (Meyer-Grass and Schneebeil 1992; Bebi et al. 2009). By experimenting with different management regimes on the DMD (Fig. 8), we concluded that Norway spruce stands could remain within a low-risk zone for as long as 60 years, provided that site index is not too high, such as most subalpine stands (e.g., 25.2 m on average for stands at elevations  $>1,700 \text{ m}$  on the Eastern Alps, data from Cantiani et al. 2000). Even for high potential productivity, the low-risk period could extend up to 30 years, which would allow for spatial planning of silvicultural interventions in avalanche-prone catchments, with a goal to maintain some proportion of Norway spruce stands in the catchment as active protection forests. Boundaries for the low-risk zone could be extended by relaxing the tree slenderness or competitive status requirements. However, this would come at the expense of individual vitality and stand-scale resistance. When the degree of tree clumping is high, it is very difficult to contrast the presence of gaps large enough to trigger potentially hazardous snow movements. Management can mitigate the tendency for large gap creation at lower elevations. For example, simulations by Cordonnier et al. (2008) suggest that by creating small gaps every 20 years, uneven-aged structure can be initiated, thereby increasing the protective function of mountain Norway spruce stands in the western Alps. In subalpine forests, which exhibit clumped spatial arrangements (Motta and Lingua 2005), stabilization of avalanche channels has to be pursued by alternative means or structures. Similar considerations

**Fig. 7** DMD and risk zones for windfirmness of Norway spruce stands. Starting stand conditions, EOR, and unmanaged stand trajectory as in working example for Fig. 6



could be made for rockfall, albeit using different thresholds on the DMD (Vacchiano et al. 2008).

#### Resistance to spruce bark beetle


In central-southern Europe, spruce bark beetle outbreaks are a part of the natural disturbance regimes of Norway spruce forests (Svoboda et al. 2012). However, mortality induced by bark beetle may severely alter structure and functionality of stands that are managed for important ecosystem services, such as protection from geological hazards (Amman 2006) or water quality (Huber 2005). Outbreaks are primarily triggered by climate and abundance of infestation source such as recent deadwood; droughts, windthrow, or pollution may decrease tree vigor and increase susceptibility, although evidence is still contradictory to this extent (Baier 1996; Dutilleul et al. 2000; Wermelinger 2004). Norway spruce trees have recently been found to be potentially more resistant to spruce bark beetle when the density of foliage or foliage packing is high (Jakuš et al. 2011), presumably as a result of the

inability of adults to reach the stem. This suggests Norway spruce trees that maintain longer crowns *throughout stand development* are more likely to resist spruce bark beetle infestation. Although the DMD was developed using stand-level data, it is relatively easy to visualize stand-density combinations necessary to maintain long live crowns. If we were to assume that full canopy closure in Norway spruce stands occurs at 25–35 % SDI (Long 1985), we would seek to maintain stands on average below that level when portrayed on the DMD. While it may be possible to enhance individual-tree growth and potentially resist the beetle under this regime, it would come at the expense of stand-level growth and would almost certainly result in low-quality logs by the EOR because of large lower branches. This shows that trade-offs associated with management goals must be considered. Fortunately, they can be simultaneously portrayed on the DMD.

An overlay of low-risk zones from Figs. 7 and 8 demonstrates potential conflicting management goals or desired conditions that cannot be simultaneously maximized. The ability of Norway spruce stands to meet various management



**Fig. 8** Low-risk zone for avalanche release hazard (slope = 35°). Low-risk boundaries express: *a* minimum basal area, *b* SDI for minimum crown ratio, *c* maximum HT<sub>100</sub>/DD ratio. *d*—red lines maximum gap size for NNI = 0.5 (clumped tree spatial pattern) and 1 (random pattern) according to Eq. 7. Starting stand conditions, EOR, and unmanaged stand trajectory as in working example for Fig. 6



objectives can be assessed on the DMD provided that associated requirements can be expressed by average (or distributional) stand parameters. Possibilities include habitat quality for ungulates (Smith and Long 1987) and birds (Shaw and Long 2007). For example, the DMD can be used to project which density regime would promote tree growth of the dominant cohort and speed up the creation of future veteran trees that will serve as habitat when alive or standing dead or to estimate the time necessary for conversion from monocultures to mixed natural forest by using the MSB to manage for time required to form stable canopy gaps.

## Conclusion

The proposed DMD represents a marked improvement in Norway spruce density management over conventional approaches, because it characterizes ecological processes that drive growth and mortality. Statistical results for the stand-scale DMD suggest it is adequately robust for use over the geographic area covered by our analysis. The

DMD allows the silviculturist to graphically display current stand conditions and project stand development after treatment with respect to density-dependent mortality and susceptibility of stand structure to natural hazards or disturbance agents. Multiple management scenarios can be simultaneously portrayed on the DMD to assess which EOR goals in terms of tree size, density, volume, and ecosystem services can be met, how much time is required to meet them, and how long they can be maintained by management.

**Acknowledgments** We acknowledge data contributors and in particular Roberta Berretti and Daniele Castagneri (Università di Torino, Italy), Momchil Panayotov (University of Forestry, Sofia, Bulgaria) Thomas Vrska (Agency for Nature Conservation and Landscape Protection of the Czech Republic), Fabio Maistrelli (Bolzano Autonomous Province, Italy), Hubert Hasenauer (University of Natural Resources and Applied Life Sciences, Vienna, Austria), IPLA SpA (regional forest inventory produced by Regione Piemonte with the support of EU structural funds), and Regione Autonoma Valle d'Aosta. Raw data were also provided by the Italian National Forest Inventory of Forests and Forest Carbon Pools (INFC 2005), carried out by the National Forest Service of Italy (CFS) and the Research Council for Agriculture—Forest Monitoring and Management

Research Unit (CRA-MPF). Support to M. Svoboda to this research came from the Czech Science Foundation (GACR P504/10/1644).

## References

- Amman M (2006) Schutzwirkung abgestorbener Bäume gegen Naturgefahren. PhD dissertation. Eidgenössische Forschungsanstalt für Wald, Schnee und Landschaft WSL, Birmensdorf
- Angelstam P, Kuuluvainen T (2004) Boreal forest disturbance regimes, successional dynamics and landscape structures—a European perspective. *Ecol Bull* 51:117–136
- Aulitzky H (1984) The microclimatic conditions in a subalpine forest as basis for the management. *GeoJournal* 8:277–281
- Baier P (1996) Defence reactions of Norway spruce (*Picea abies* Karst.) to controlled attacks of *Ips typographus* (L.) (Col., Scolytidae) in relation to tree parameters. *J Appl Entomol* 120:587–593
- Bebi P, Kienast F, Schönenberger W (2001) Assessing structures in mountain forests as a basis for investigating the forests' dynamics and protective functions. *For Ecol Manag* 145:3–14
- Bebi P, Kulakowski D, Rixen C (2009) Snow avalanche disturbances in forest ecosystems—State of research and implications for management. *For Ecol Manag* 257:1883–1892
- Berretti R, Motta R (2005) Ungulati selvatici e foresta. Parco Naturale di Paneveggio—Pale di S Martino, Trento
- Berretti R, Caffo L, Camerano P, De Ferrari F, Domaine A, Dotta A, Gottero F, Haudemand JC, Letey C, Meloni F, Motta R, Terzuolo PG (2006) Selvicoltura nelle foreste di protezione: Esperienze e indirizzi gestionali in Piemonte e Valle d'Aosta. Compagnia delle Foreste, Arezzo
- Bi H, Turvey ND (1997) A method of selecting data points for fitting the maximum biomass–density line for stand undergoing self-thinning. *Aust J Ecol* 22:356–359
- Buford MA, Burkhardt HE (1987) Genetic improvement effects on growth and yield of loblolly pine plantations. *For Sci* 33:704–724
- Cantiani MG, Floris A, Tabacchi G (2000) Yield features of high mountain and subalpine Spruce forests in Val di Fiemme (Trentino, Italy). Comunicazione di Ricerca dell'Istituto Sperimentale per l'Assessment Forestale e per l'Alpicoltura 3:3–21
- Castagneri D, Vacchiano G, Lingua E, Motta R (2008) Analysis of intraspecific competition in two subalpine Norway spruce (*Picea abies* (L.) Karst.) stands in Paneveggio (Trento, Italy). *For Ecol Manag* 255:651–659
- Castagneri D, Garbarino M, Berretti R, Motta R (2010) Site and stand effects on coarse woody debris in montane mixed forests of Eastern Italian Alps. *For Ecol Manag* 260:1592–1598
- Castedo-Dorado F, Crecente-Campo F, Alvarez-Alvarez P, Barrio-Anta M (2009) Development of a stand density management diagram for radiata pine stands including assessment of stand stability. *Forestry* 82:1–16
- Cenușă R (1992) Cercetări asupra structurii volumului ecologic și succesiunii ecosistemelor forestiere de limită altitudinală din Carpații Nordici (Călimani și Giumalău). PhD Dissertation, Academia de Științe Agricole și Silvicultură, București
- Charru M, Seynave I, Morneau F, Bontemps JD (2011) Significant differences and curvilinearity of the self-thinning relationship of eleven species based on forest inventory data. *Ann For Sci* 69:195–205
- Čížková P, Svoboda M, Křenová Z (2011) Natural regeneration of acidophilous spruce mountain forests in non-intervention management areas of Šumava National Park—the first results of the Biomonitoring project. *Silva Gabreta* 17:19–35
- Clark PJ, Evans FC (1954) Distance to nearest neighbor as a measure of spatial relationships in populations. *Ecology* 35:445–453
- Cordonnier T, Courbaud B, Berger F, Franc A (2008) Permanence of resilience and protection efficiency in mountain Norway spruce forest stands: a simulation study. *For Ecol Manag* 256:347–354
- Curtis R (2010) Effect of diameter limits and stand structure on relative density indices: a case study. *West J Appl For* 25:169–175
- Dobbertin M (2002) Influence of stand structure and site factors on wind damage comparing the storms Vivian and Lothar. *For Snow Landsc Res* 77:187–205
- Drew TJ, Flewelling JW (1979) Stand density management: an alternative approach and its application to Douglas-fir plantations. *For Sci* 25:518–532
- Ducey MJ (2009) The ratio of additive and traditional stand density indices. *West J Appl For* 24:5–10
- Dutilleul P, Nef L, Frigon D (2000) Assessment of site characteristics as predictors of the vulnerability of Norway spruce (*Picea abies* Karst.) stands to attack by *Ips typographus* L. (Col., Scolytidae). *J Appl Entomol* 124:1–5
- Fisher A, Lindner M, Abs C, Lasch P (2002) Vegetation dynamics in Central European forest ecosystems (near natural as well as managed) after storm events. *Folia Geobot* 37:17–21
- Gamborg C, Larsen JB (2003) Back to nature—a sustainable future for forestry? *For Ecol Manag* 179:559–571
- Gauquelin X, Courbaud B (eds) (2006) Guide de sylviculture des forêts de montagne—Alpes du Nord françaises. Cemagref et Office National des Forêts, Grenoble
- Hansen J, Spiecker H (2004) Conversion of Norway spruce (*Picea abies* [L.] Karst.) forests in Europe. In: Stanturf JA, Madsen P (eds) Restoring temperate and boreal forested restoration of boreal and temperate forests. CRC Press, Boca Raton, pp 339–347
- Huber C (2005) Long lasting nitrate leaching after bark beetle attack in the highlands of the Bavarian Forest National Park. *J Environ Qual* 34:1772–1779
- Hynynen J, Ojansuu R (2003) Impact of plot size on individual-tree competition measures for growth and yield simulators. *Can J For Res* 33:455–465
- INFC (2005) Linee generali del progetto per il secondo inventario forestale nazionale italiano. CRA-ISAFA, MiPAF—Direzione Generale per le Risorse Forestali Montane e Idriche, Corpo Forestale dello Stato, Trento
- INFC (2006) Procedure di posizionamento e di rilievo degli attributi di terza fase. CRA-ISAFA, MiPAF—Direzione Generale per le Risorse Forestali Montane e Idriche, Corpo Forestale dello Stato, Trento
- IPLA (2003) Manuale dei rilievi inventariali di campagna. Regione Piemonte, Torino
- Jack SB, Long JN (1996) Linkages between silviculture and ecology: an analysis of density management diagrams. *For Ecol Manag* 86:205–220
- Jakuš R, Edwards-Jonášová M, Cudlín P, Blaženec M, Ježík M, Havlíček F, Moravec I (2011) Characteristics of Norway spruce trees (*Picea abies* Karst.) surviving a Spruce bark beetle (*Ips typographus* L.) outbreak. *Trees* 25:965–973
- Klopčič M, Poljanec A, Gartner A, Boncina A (2009) Factors related to natural disturbances in mountain Norway spruce (*Picea abies*) forests. *Ecoscience* 16:48–57
- Körner C (2003) Alpine plant life: functional plant ecology of high mountain ecosystems. Springer, Berlin
- Krumm F, Kulakowski D, Spiecker H, Duc P, Bebi P (2011) Stand development of Norway spruce dominated subalpine forests of the Swiss Alps. *For Ecol Manag* 262:620–628
- Krumm F, Kulakowski D, Risch AC, Spiecker H, Bebi P (2012) Stem exclusion and mortality in unmanaged subalpine forests of the Swiss Alps. *Eur J For Res* 131:1571–1583

- Kulakowski D, Bebi P (2004) Range of variability of unmanaged subalpine forests. *Forum für Wissen* 2004:47–54
- Long JN (1985) A practical approach to density management. *For Chron* 61:23–27
- DeRose RJ, Shaw JD, Vacchiano G, Long JN (2008) Improving longleaf pine mortality predictions in the Southern Variant of the Forest Vegetation Simulator. In: Havis RN, Crookston NL (eds) 2008 Third Forest Vegetation Simulator Conference; Fort Collins, February 13–15, 2007; Proceedings RMRS–P–54 Fort Collins, CO: USDA Forest Service, Rocky Mountain Research Station: pp 160–166
- Long JN, Daniel TW (1990) Assessment of growing stock in uneven aged stands. *West J Appl For* 5:93–96
- Long JN, Shaw JD (2005) A density management diagram for even-aged ponderosa pine stands. *West J Appl For* 20:205–215
- Mayer H, Ott E (1991) *Gebirgswaldbau—Schutzwaldpflege*. Gustav Fischer, Stuttgart
- Meyer-Grass M, Schneebeli M (1992) Die Abhängigkeit der Waldlawinen von Standorts-, Bestandes- und Schneebedingungen. Internationales Symposium Interpraevent 1992-Bern, Tagungspublikation, Band 2: 443–455
- Mitchell S (2000) Forest health: preliminary interpretations for wind damage. BC Ministry of Forests, Forest Practices Branch, Victoria
- Monserud RA, Ledermann T, Sterba H (2005) Are self-thinning constraints needed in a tree-specific mortality model? *For Sci* 50:848–858
- Moser M (1991) *Taxationshilfen für Südtirol*. Dissertation, Universität für Bodenkultur, Wien
- Motta R (2002) Old-growth forests and silviculture in the Italian Alps: the case-study of the strict reserve of Paneveggio (TN). *Plant Biosystems* 136:223–232
- Motta R, Haudemand JC (2000) Protective forests and silvicultural stability—an example of planning in the Aosta Valley. *Mount Res Devel* 20:180–187
- Motta R, Lingua E (2005) Human impact on size, age, and spatial structure in a mixed European larch and Swiss stone pine forest in the Western Italian Alps. *Can J For Res* 35:1809–1820
- Motta R, Berretti R, Lingua E, Piussi P (2006) Coarse woody debris, forest structure and regeneration in the Valbona Forest Reserve, Paneveggio, Italian Alps. *For Ecol Manag* 235:155–163
- Nascimbene J, Marini L, Motta R, Nimis PL (2009) Influence of tree age, tree size and crown structure on lichen communities in mature Alpine spruce forests. *Biodiv Conserv* 18:1519–1522
- Panayotov M, Kulakowski D, Laranjeiro DS, Bebi P (2011) Wind disturbances shape old Norway spruce-dominated forest in Bulgaria. *For Ecol Manag* 262:470–481
- Parviainen J, Kassiomis K, Bücking W, Hochbichler E, Päivinen R, Little D (2000) Final report summary: mission, goal, outputs, linkages, recommendations and partners. In: Commission European (ed) EUR 19550 – COST action E4—forest reserves research network. Office for Official Publication of the European Communities, Luxembourg, pp 9–38
- Pretzsch H (2005) Stand density and growth of Norway spruce (*Picea abies* (L.) Karst) and European beech (*Fagus sylvatica* L.): evidence from long-term experimental plots. *Eur J For Res* 124:193–205
- Pretzsch H (2006) Species-specific allometric scaling under self-thinning. Evidence from long-term plots in forest stands. *Oecologia* 146:572–583
- Pretzsch H, Biber P (2005) A re-evaluation of Reineke's rule and stand density index. *For Sci* 51:304–320
- Pretzsch H, Grote R, Reineking B, Rötzer T, Seifert S (2008) Models for forest ecosystem management: a European perspective. *Ann Bot* 101(8):1065–1087
- Reineke LH (1933) Perfecting a stand-density index for even-aged forests. *J Agric Res* 46:627–638
- Riou-Nivert P (2001) Facteurs de stabilité des peuplements et gestion de l'équilibre. *Forêt Entreprise* 139:17–25
- Rottmann M (1986) Wind- und Sturmschäden im Wald. Beiträge zur Beurteilung der Bruchgefährdung, zur Schadensvermeidung und zur Behandlung sturmgeschädigter Nadelholzbestände. Sauerländer, Frankfurt am Main
- Schelhaas MJ, Nabuurs GJ, Schuck A (2003) Natural disturbances in the European forests in the 19th and 20th centuries. *Global Change Biol* 9:1620–1633
- Schlyter P, Stjernquist I, Barring L, Jönsson AM, Nilsson C (2006) Assessment of the impacts of climate change and weather extremes on boreal forests in northern Europe, focusing on Norway spruce. *Clim Res* 31:75–84
- Schmidt M, Kandler G (2009) An analysis of Norway spruce stem quality in Baden-Württemberg: results from the second German national forest inventory. *Eur J For Res* 128:515–529
- Schmidt-Vogt, H (1977) Die Fichte. Band I Taxonomie – Verbreitung – Morphologie – Ökologie – Waldgesellschaften. Paul Parey, Hamburg und Berlin
- Schmidt-Vogt H, Wütherich G, Deichner P (1987) Untersuchungen zur Sturmstabilität von Fichten und Tannen in Fichten-Tannen-Mischbeständen auf verschiedenen Standorten Süddeutschlands. *Allgemeine Forst- und Jagdzeitung* 158:42–50
- Schönenberger W (2001) Cluster afforestation for creating diverse mountain forest structures—a review. *For Ecol Manag* 145:121–128
- Schütz JP, Zingg A (2010) Improving estimations of maximal stand density by combining Reineke's size-density rule and the yield level, using the example of spruce (*Picea abies* (L.) Karst) and European Beech (*Fagus sylvatica* L.). *Ann For Sci* 67:507
- Schütz JP, Götz M, Schmid W, Mandallaz D (2006) Vulnerability of spruce (*Picea abies*) and beech (*Fagus sylvatica*) forest stands to storms and consequences for silviculture. *Eur J For Res* 125:291–302
- Seidl R, Schelhaas MJ, Lexer MJ (2011) Unraveling the drivers of intensifying forest disturbance regimes in Europe. *Global Change Biol* 17:2842–2852
- Shaw JD (2000) Application of stand density index to irregularly structured stands. *West J Appl For* 15:40–42
- Shaw JD, Long JN (2007) A density management diagram for longleaf pine stands with application to red-cockaded woodpecker habitat. *South J Appl For* 31:28–38
- Shorohova E, Fedorchuk V, Kuznetsova M, Shvedova O (2008) Wind-induced successional changes in pristine boreal *Picea abies* forest stands: evidence from long-term permanent plot records. *Forestry* 81:335–359
- Shorohova E, Kuuluvainen T, Kangur A, Jögi K (2009) Natural stand structure, disturbance regimes, and successional dynamics in the Eurasian boreal forest: a review with special reference to Russian studies. *Ann For Sci* 66:201
- Skrøppa T (2003) Euforgen technical guidelines for genetic conservation and use for Norway spruce (*Picea abies*). International Plant Genetic Resources Institute, Rome
- Smith FW, Long JN (1987) Elk hiding and thermal cover guidelines in the context of lodgepole pine stand density. *West J Appl For* 2:6–10
- Sterba H (1976) Die Funktionsschemata der vier Fichtenertragstafeln. *Centralblatt für das gesamte Forstwesen* 93:102–112
- Sterba H (1981) Natürlicher Bestockungsgrad und Reinekes SDI. *Centralblatt für das gesamte Forstwesen* 98:101–116
- Sterba H (1987) Estimating potential density from thinning experiments and inventory data. *For Sci* 33:1022–1034
- Svoboda M, Fraver S, Janda P, Bace R, Zenahlikova J (2010) Natural development and regeneration of a Central European montane spruce forest. *For Ecol Manag* 260:707–714
- Svoboda M, Janda P, Nagel TA, Fraver S, Rejzek J, Bace R (2012) Disturbance history of an old-growth sub-alpine *Picea abies*

- 888 stand in the Bohemian Forest, Czech Republic. J Veg Sci 903  
 889 23:86–97 904  
 890 Thomasius H (1980) Wissenschaftliche Grundlagen der Rahmenrich- 905  
 891 tlinie zur Behandlung bruchgeschädigter Fichten- und Kie- 906  
 892 fernbestände. Sozialistische Forstwirtschaft 30:364–373 907  
 893 Thomasius H (1988) Stabilität natürlicher und künstlicher Waldöko- 908  
 894 systeme sowie deren Beeinflussbarkeit durch forstwirtschaftliche 909  
 895 Massnahmen (Teil I and II). Allg For Z 43(1037–1043):1064–1068 910  
 896 Vacchiano G, Motta R, Long JN, Shaw JD (2008) A density 911  
 897 management diagram for Scots pine (*Pinus sylvestris* L): a tool 912  
 898 for assessing the forest's protective effect. For Ecol Manag 913  
 899 255:2542–2554 914  
 900 Vidal C, Bélouard T, Hervé JC, Robert N, Wolsack J (2007) A new 915  
 901 flexible forest inventory in France. In: McRoberts RE, Reams 916  
 902 GA, Van Deusen PC, McWilliams WH (eds), Proceedings of the 917  
 7th Annual Forest Inventory and Analysis Symposium, Portland 918  
 3–6 Oct 2005, General Technical Report WO-77, USDA Forest  
 Service, pp 67–73  
 Weiner J (2004) Allocation, plasticity and allometry in plants. Perspect Plant Ecol Evol Syst 6:207–215  
 Wermelinger B (2004) Ecology and management of the spruce bark beetle *Ips typographus*—a review of recent research. For Ecol Manag 202:67–82  
 Yoda K, Kira T, Ogawa H, Hozumi K (1963) Self-thinning in overcrowded pure stands under cultivated and natural conditions. J Biol Osaka City Univ 14:107–129  
 Zajaczkowski J (1991) Odporność lasu na szkodliwe działanie wiatru i śniegu. Wydawnictwo Świat, Warsaw  
 Zeide B (1985) Tolerance and self-tolerance of trees. For Ecol Manag 13:149–166